



(11) Publication number:

**0 101 124
A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 83201137.3

(22) Date of filing: 01.08.83

(51) Int. Cl.³: **C 07 C 67/475**
C 07 C 69/734, C 07 C 51/00
C 07 C 59/64, C 07 D 333/22
C 07 C 57/30

(30) Priority: 06.08.82 IT 2276082
07.03.83 IT 1993083

(43) Date of publication of application:
22.02.84 Bulletin 84/8

(84) Designated Contracting States:
AT BE CH DE FR GB IT LU NL SE

(71) Applicant: ZAMBON S.p.A.
Via Cappuccini, 40
I-36100 Vicenza(IT)

(72) Inventor: Giordano, Claudio
Via Canove Nuove, 23
I-36100 Vicenza(IT)

(72) Inventor: Castaldi, Graziano
Via Livia Gallina, 5
Briona (Novara)(IT)

(74) Representative: Marchi, Massimo et al,
c/o CON.P.I. Corso Buenos Aires, 64
I-20124 Milano(IT)

(54) Process for preparing alpha arylalkanoic acids.

(57) Rearrangement of alpha-halo-alkylarylketals in neutral or weakly alkaline conditions and in the presence of a polar-protic medium and subsequent hydrolysis of the thus obtained esters, in the same reaction medium, to afford the corresponding alpha-arylalkanoic acids or their salts which are particularly useful as anti-inflammatory, analgesic and antipyretic agents.

EP 0 101 124 A1

"PROCESS FOR PREPARING ALPHA ARYLALKANOIC ACIDS"

* * * * *

5 This invention relates to a process for preparing an alpha-aryl
alkanoic acid or a salt thereof which comprises the rearrangement
of an alpha-halo-alkylarylketal in neutral or weakly alkaline con-
ditions and in the presence of a polar-protic medium and the sub-
sequent hydrolysis of the thus obtained ester in the same reaction
medium to afford the corresponding alpha-arylalkanoic acid.

10 The European Patent Application 34871 describes a process for
preparing esters of alpha-arylalkanoic acids via rearrangement of
alpha-halo-alkylarylketals in the presence of a Lewis acid. The
disadvantages of this process are that the reaction must be car-
ried out in an anhydrous medium and that the more effective Lewis
acids are the salts of toxic heavy metals; consequently there is a
15 need for a careful purification when the final product is intended
for pharmaceutical use. This purification step may be carried out
sucessfully by isolating the ester but this impairs the compaction
of both the rearrangement and the hydrolysis steps.

20 Furthermore the Lewis acids incline to ineract with the oxygen
atoms of the ketal group and, consequently, to form by-products
whose amount ranges in accordance with the nature of the alpha-ha-
lo-alkylarylketal and of the Lewis acid which are used.

25 The European Application No. 48.136 describes a process for pre-
paring alpha-arylalkanoic acids wherein an alpha-sulfonyloxyketal
undergoes hydrolysis. This process lays on the prejudice that the
halogen atom is not sufficiently labile to promote the rearrange-
ment in the absence of a catalyst having affinity for halogen
whereas it was well known that the sulfonyloxy-groups like the to-
syloxy and mesyloxy are sufficiently labile to disassociate from
30 the substrate upon contact with a protic-polar medium.

Despite the apparent easy of the last step this process is complex and cumbersome owing to the fact that at first an alpha-halo-ketone is reacted with an alkali metal alkoxide to afford an alpha-hydroxyketal which is then treated with an O-sulfonylating agent to form the corresponding alpha-sulfonyloxyketal and finally the alpha-sulfonyloxyketal is submitted to hydrolysis. The number of steps involved and the difficulties inherent in carrying out on a large scale some of the steps leave room for a more simple and economical process.

Now it has been surprisingly found that the alpha-haloalkyl aryl ketals undergo rearrangement in neutral or weakly alkaline conditions and in the presence of a protic-polar medium thus avoiding the detour depicted by the European Patent Application 48.136 as well as disadvantages inherent in the process described by the European Patent Application 34.871.

More particularly the process according to this invention comprises the rearrangement of an alpha-haloalkylarylketal in neutral or weakly alkaline conditions and in the presence of a polar-protic medium and the subsequent hydrolysis of the thus obtained ester in the same reaction medium to afford the corresponding alpha-aryl alkanolic acid or a salt thereof.

The neutral or weakly alkaline condition is obtained by adding buffers or weak bases either organic or inorganic or mixture thereof. Examples of typical compounds are the aliphatic and the aromatic tertiary amines and the alkali and alkaline earth metals salts of organic and inorganic acids such as sodium bicarbonate, calcium carbonate, potassium acetate, triethylamine, methylpiperidine, methyl pyrrolidine and dimethylaniline.

The protic-polar media according to this invention are water, alcohols and mixtures thereof. Examples of alcohols include C₁-C₁₂

primary, secondary and tertiary alcohols and C_2-C_{12} polyhydric alcohols such as methanol, ethanol, butanol, isobutanol, sec butanol, 2-decanol, allyl alcohol, ethylene glycol, 1,2-propanediol, 1,3-propanediol, 1,10-decanediol, cis-2-buten-1,4-diol and the like.

Preferably the choice of the protic medium will depend from its lipophilic and hydrophilic properties as well as from the nature of the alpha-halo-alkylaryl ketal.

As a rule a decrease in the concentration of the hydroxy- groups such as in the alcohol and diols having a long aliphatic chain, increases the lipophilic property of the protic medium and in its turn promotes the solubilization of the low-polar ketals but reduces their reaction rate which, on the contrary is favored by strongly hydrophilic protic media. Of course, the reaction rate is increased also by an increase in the reaction temperature.


Usually the reaction time is comprised between few minutes and about twelve hours when a ketal endowed with a normal reactivity is treated at a temperature from 80° C to 200° C.

When the rearrangement is complete the pH of the reaction mixture is made acid or basic for hydrolyzing the ester. Preferably the pH is increased by addition of a base. Example of suitable bases include alkali and alkaline earth metal hydroxides and salts. Usually this step is complete in a time period comprised from few minutes and some hours at a temperature from 20° to 100°C. It is so obtained a salt of an alpha-arylalkanoic acid which is then treated, when desired, with an acid to afford the free alpha-arylalkanoic acid.

Examples of arylalkanoic acids which may be prepared with the process of this invention include aclofenac, benoxaprofen, caroprofen, diclofenac, fenclofenac, fenoprofen, fentiazac, flurbiprofen,

indoprofen, ibuprofen, isoprofen, ketoprofen, naproxen, piroprofen, suprofen, tolmetin, xenbucin and the like.

The preferred ketals used as starting products according to this invention have the following formula



wherein A_x is an aromatic ring selected from the group comprising

- (a) a phenyl ring substituted by one or two substituents selected from the group comprising halogen, 1-6 C alkyl, 1-4 C alkoxy, 2-4 C alkenyloxy, phenyl, phenoxy, dichlorophenoxy, dichloroanilino, benzoyl, indolinyll, dihydropyrrolyl, thenoyl (b) a naphthyl ring substituted by one or two substituents selected from the group comprising halogen and 1-4 C alkoxy
- (c) a pyrrolyl ring substituted by one or two radicals selected from the group comprising 1-4 C alkyl and alkyl_{1-4 C}-phenyl, (d) chloro-carbazolyl, (e) benzoxazolyl substituted by one chlorophenyl radical,
- (f) thiazolyl substituted by one or two radical selected from the group comprising phenyl and chloro phenyl, and (g) thienyl,

R' and R'' , independently from each other, are a saturated or unsaturated, straight or branched alkyl radical having from 1 to 12 C atoms or, when taken together, are a saturated or unsaturated straight or branched alkylene radical having from 2 to 12 C atoms which completes an alicyclic ring having from 5 to 7 members,

X is halogen;

R''' is hydrogen, alkyl having from 1 to 6 C atoms
or cycloalkyl having from 1 to 7 C atoms.

The alpha-halo-alkylarylketals may be prepared according to known methods.

5 Examples of methods suitable for preparing alpha-halo-alkylaryl
ketals contemplate the use of an acid catalyst such as p-toluene-
sulfonic acid (J. Org. Chem. 21, 1366 (1956); ibidem, 22, 662,
(1957); Synthesis 23, (1974), active montmorillonite (Bull. Soc.
Chim. France, 2558, (1975)), BF₃ etherate (Bull. Soc. Chim. France
10 1763 (1975)) and citric acid (U.S. Patent 3.936.398). An improve-
ment to the last method contemplates the use of citric acid in the
presence of hydroquinone which acts as polymerization inhibitor
(Bull. Soc. Chim. France, 1973 (1975)).

15 The water generated in the course of the reaction is removed by
azeotropic distillation with suitable solvents such as benzene,
toluene, cyclohexane and the like or by means of a dehydrating a-
gent such as anhydrous CuSO₄, a trialkylorthoformate, molecular
sieves and the like (Synthesis, 501 (1981)).

20 Other known methods for preparing the ketals are the trans-ke-
talization, the reaction of a ketone with an alcohol in the pre-
sence of a ketal such as 2,2-dimethoxypropane and an acid catalyst
(J. Org. Chem. 25, 521 (1960)) and the reaction of an alcohol with a
suitable enol-ether in the presence of an acid catalyst (Bull. Soc.
Chim. France, 264 (1979)).

25 Examples of known alpha-halo-alkylarylketals include:

- 2-bromo-1,1-dimethoxy-1-(6'-methoxy-2'-naphthyl)-propane,
- 2-bromo-1,1-dimethoxy-1-(4'-isobutylphenyl)-propane,
- 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxane, and
- 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxolane.

30 Examples of not-yet described alpha-halo-alkylarylketals which

are prepared according to known techniques are:

2-bromo-1,1-dimethoxy-1-(4'-methoxyphenyl)-propane

¹H-NMR (60 MHz) (CDCl₃ - TMS) delta (ppm): 1.45 (d, 3H); 3.15 (s, 3H); 3.8 (s, 3H); 4.5 (q, 1H); 6.85-7.6 (m, 4H).

5 2-bromo-1,1-dimethoxy-1-(4'-isobutylphenyl)-propane

¹H-NMR (60 MHz) (CDCl₃ - TMS) delta (ppm): 0.9 (d, 6H); 1.5 (d, 3H); 1.7-2.2 (m, 1H); 2.6 (d, 2H); 3.2 (s, 3H); 3.4 (s, 3H); 4.5 (q, 1H); 7.1-7.6 (m, 4H).

10 2-(1'-bromoethyl)-5,5-dimethyl-2-(6'-methoxy-2'-naphthyl)-1,3-dioxane, m.p. 89-90° C.

2-(1'-bromoethyl)-2-(4'-methoxyphenyl)-1,3-dioxane

¹H-NMR (60 MHz) (CDCl₃ - TMS) delta (ppm): 1.65 (d, 3H), 3.6-4 (m, 6H); 3.85 (s, 3H); 4.1 (q, 1H); 7-7.7 (m, 4H).

15 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxane

¹H-NMR (60 MHz) (CDCl₃ - TMS) delta (ppm): 1.20 (m, 2H); 1.68 (d, 3H); 3.90 (m, 4H); 3.96 (s, 3H); 4.30 (q, 1H); 7.12-7.98 (m, 6H).

2-bromoethyl)-5-(2'-butyl)-5-methyl-2-(4'-methoxyphenyl)-1,3-dioxane,

20 2-(1'-bromoethyl)-5,5-dimethyl-2-(4'-isobutylphenyl)-2,3-dioxane, b.p. 135-138° C (0.6 mm Hg)

2-(1'-bromoethyl)-5,5-dimethyl-2-(5'-bromo-6'-methoxy-2'-naphthyl)-1,3-dioxane, m.p. 143-145° C

2-(1'-bromoethyl)-5,5-dimethyl-2-(2'-thienyl)-1,3-dioxane, b.p. 78-80° C (0.5 mm Hg)

25 2-(1'-bromoethyl)-2-(4'-methoxyphenyl)-1,3-dioxolane

¹H-NMR (60 MHz) (CDCl₃ - TMS) (ppm): 1.6 (d, 3H), 3.8 (s, 3H); 3.8-4.2 (m, 4H); 4.4 (q, 1H); 6.85 - 7.6 (m, 4H)

30 Particularly useful new alpha-halo-alkylarylketals of formula I are those where R' and R" together are a straight or branched unsaturated alkylidene radical which forms a ring having 7 members.

They are described by the Italian Patent Application No. 19930 A/83 filed on March 7, 1983 and are an object of the Patent Application No. filed on

5 The following example are given to illustrate this invention without limiting it in any way.

EXAMPLE 1

2-(6'-methoxy-2'-naphthyl)-propionic acid

10 a) A mixture of 2-bromo-1,1-dimethoxy-2-(6'-methoxy-2'-naphthyl)-propane (3.39 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and 1,10-decane diol (5 g) is heated to 170° C for six hours.

15 After having reduced the temperature to 70° C and added 30% potassium hydroxide, the reaction mixture is heated to 100° C for five hours. The mixture is then cooled to room temperature, poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml).

20 The aqueous layer is acidified with concentrated hydrochloric acid to afford a precipitate consisting of 2-(6'-methoxy-2'-naphthyl)-propionic acid (1.38 g; 0.006 mol). Yield, 60%; m.p. 155-156° C.

b) By substituting ethylene glycol (40 ml) for 1,10-decanediol in the procedure of Example 1a and otherwise heating the reaction mixture to 125° C for 8 hours, there is obtained the title product (1.9 g; 0.0082 mol). Yield, 82%; m.p. 155-157° C.

25 c) By substituting allyl alcohol (50 ml) for 1,10-decanediol in the procedure of example 1a and otherwise refluxing the reaction mixture for 48 hours, there is obtained the title product (1.2 g; 0.0052 mol). Yield, 52%; m.p. 155-156° C.

30 d) A mixture of 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxolane (3.37 g; 0.01 mol), potassium acetate (1.2 g; 0.012

mol) and ethylene glycol (50 ml) is heated to 125° C for 16 hours.

By following the procedure of Example 1a, there is obtained the title product (1.61 g; 0.007 mol). Yield, 70%; m.p. 155-156° C.

- 5 e) A mixture of 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxolane (2.68 g; 0.008 mol), disodium phosphate (1.36 g; 0.0095 mol), monopotassium phosphate (1.44 g; 0.01 mol) and ethylene glycol (40 ml) is heated to 125° C for 16 hours.

10 By following the procedure of Example 1a, there is obtained the title product (1.25; 0.005 mol). Yield, 62,5%; m.p. 155-156° C.

- f) A mixture of 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxolane (13.48 g; 0.042 mol), potassium bicarbonate (6 g; 0.06 mol) and ethylene glycol (200 ml) is heated to 125° C for 18 hours.

15 By following the procedure of Example 1a, there is obtained the title product (6.91 g; 0.03 mol). Yield, 71,5%; m.p. 155-156° C.

- g) A mixture of 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxane (3.51 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and 1,3-propanediol (50 ml) is heated to 125° C for 8 hours.

20 By following the procedure of Example 1a, there is obtained the title product (2.25 g; 0.0098 mol). Yield, 98%; m.p. 155-156° C.

- h) A mixture of 2-(1'-bromoethyl)-5,5-dimethyl-2-(6'-methoxy-2'-naphthyl)-1,3-dioxane (3.79 g; 0,01 mol), potassium acetate (1.2 g; 0,012 mol) and ethylene glycol (40 ml) is heated to 125° C for 4 hours.

25 By following the procedure of Example 1a, there is obtained the title product (2.1 g; 0.0091 mol). Yield, 91%; m.p. 154-155° C.

- i) A mixture of 2-(6'-methoxy-2'-naphthyl)-2-(1'-bromoethyl)-4,7-dihydro-1,3-dioxepine (1.09 g; 0.003 mol), cis-2-butene-1,4-diol (10 ml) and potassium acetate (0.40 g; 0.004 mol) is heated to
- 30

125° C for 1 hour.

By following the procedure of Example 1a, there is obtained the title product (0.6 g; 0.0026 mol). Yield, 87%; m.p. 155-156° C.

- 5 1) A mixture of 2-(1'-bromoethyl)-2-(6'-methoxy-2'-naphthyl)-1,3-dioxolane (3.37 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and 2,2-dimethyl-1,3-propanediol (30 g) is heated to 150° C for 24 hours.

By following the procedure of Example 1a, there is obtained the title product (1.61 g; 0.007 mol). Yield, 70%; m.p. 155-156° C.

10 EXAMPLE 2

2-(4'-methoxyphenyl)-propionic acid

- 15 a) A mixture of 2-bromo-1,1-dimethoxy-1-(4'-methoxyphenyl)-propane (2.89 g; 0.01 mol), calcium carbonate (1 g; 0.01 mol), methanol (7 ml) and water (3 ml) is refluxed for 35 hours. The reaction mixture is then cooled to 25° C and worked up as described in Example 1a, there is so obtained the title product (1.69 g; 0.0094 mol). Yield, 94%; m.p. 55° C.

- 20 b) A mixture of 2-bromo-1,1-dimethoxy-1-(4'-methoxyphenyl)-propane (2.89 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol), 1,2-propanediol (16 ml) and water (8 ml) is heated to 95° C for 3 hours.

By following the procedure of Example 1a, there is obtained the title product (1.71 g; 0.0095 mol). Yield, 95%; m.p. 57° C.

- 25 c) A mixture of 2-bromo-1,1-dimethoxy-1-(4'-methoxyphenyl)-propane (2.89 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and n-butanol (40 ml) is heated to 115° C for 32 hours.

By following the procedure of Example 1a, there is obtained the title product (1.60 g; 0.0089 mol). Yield, 89%; m.p. 56-57° C.

- 30 d) A mixture of 2-(1-bromoethyl)-2-(4'-methoxyphenyl)-1,3-dioxane (3.01 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and

ethylene glycol (40 ml) is heated to 115° C for 4 hours.

By following the procedure of Example 1a, there is obtained the title product (1.70 g; 0.0094 mol). Yield, 94%; m.p. 56-57° C.

5 e) A mixture of 2-(1-bromoethyl)-2-(4'-methoxyphenyl)-1,3-dioxane (3.01 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and 1,3-propanediol (40 ml) is heated to 115° C for 4 hours.

By following the procedure of Example 1a, there is obtained the title product (1.73 g; 0.0096 mol). Yield, 96%; m.p. 57° C.

10 f) A mixture of 2-(1'-bromoethyl)-2-(4'-methoxyphenyl)-1,3-dioxolane (2.87 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and ethylene glycol (40 ml) is heated to 115° C for 15 hours.

By following the procedure of Example 1a, there is obtained the title product (1.44 g; 0.008 mol). Yield, 80%; m.p. 57° C.

15 g) A mixture of 2-(1-bromoethyl)-5-(2'-butyl)-5-methyl-2-(4'-methoxyphenyl)-1,3-dioxane (3.68 g; 0.01 mol), ethylene glycol (40 ml) and potassium acetate (1.2 g; 0.012 mol) is heated to 125° C for 4 hours.

By following the procedure of Example 1a, there is obtained the title product (1.7 g; 0.0094 mol). Yield, 94%; m.p. 56° C.

20 h) A mixture of 2-(1'-bromoethyl)-2-(4'-methoxy)-4,7-dihydro-1,3-dioxepine (3.13 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and ethylene glycol (40 ml) is heated to 115° C for 1 hour.

By following the procedure of Example 1a, there is obtained the title product (1.52 g; 0.0085 mol). Yield, 85%; m.p. 57° C

25 i) By substituting cis-2-butene-1,4-diol (40 ml) for ethylene glycol in Example 2h, and otherwise heating the reaction mixture to 125° C for 3 hours and then following the procedure of Example 1a, there is obtained the title product (1.52 g; 0.0085 mol). Yield, 85%; m.p. 55-57° C.

2-(4'-isobutylphenyl)-propionic acid

5 a) A mixture of 2-bromo-1,1-dimethoxy-1-(4'-isobutylphenyl)-propane (3.15 g; 0.01 mol), potassium acetate (12 g; 0.012 mol), n-butanol (12 ml), 1,2-propanediol (16 ml) and water is heated to 95° C for 67 hours.

By following the procedure of Example 1a, there is obtained the title product (1.75 g; 0.0085 mol). Yield, 85%; m.p. 76-77° C.

10 b) By substituting ethylene glycol (100 ml) for n-butanol, 1,2-propanediol and water in the procedure of Example 3a, and otherwise heating the reaction mixture to 125°C for 16 hours and then following the procedure of Example 1a, there is obtained the title product (1.64 g; 0.008 mol). Yield, 80%; m.p. 76-77° C.

15 c) A mixture of 2-(1'-bromoethyl)-5,5-dimethyl-2-(4'-isobutylphenyl)-1,3-dioxane (3.55 g; 0.01 mol), potassium acetate (1.29 g; 0.012 mol) and cis-2-butene-1,4-diol (40 ml) is heated to 125° C for 3 hours.

By following the procedure of Example 1a, there is obtained the title product (1.93 g; 0.0093 mol). Yield, 93%; m.p. 77° C.

20 d) A mixture of 2-(1'-bromoethyl)-2-(4'-isobutylphenyl)-4,7-dihydro-1,3-dioxepine (3.39 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and ethylene glycol (50 ml) is heated to 115° C for 25 hours.

By following the procedure of Example 1a, there is obtained the title compound (1.89 g; 0.0092 mol). Yield, 92%; m.p. 77° C.

25 e) By substituting cis-2-butene-1,4-diol (40 ml) for ethylene glycol in the procedure of Example 3d, and otherwise heating the reaction mixture to 125° C for 3 hours and then following the procedure of Example 1a, there is obtained the title product (1.85 g; 0.009 mol). Yield, 90%, m.p. 76-77° C.

f) A mixture of 2-(1'-bromoethyl)-5,5-dimethyl-2-(4'-isobutyl)-1,3-dioxane (3 55 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and 2,2-dimethyl-1,3-propanediol (30 g) is heated to 150° C for 24 hours.

5 By following the procedure of Example 1a, there is obtained the title product (1.4 g; 0.007 mol). Yield, 70%; m.p. 76-77° C.

g) A mixture of 2-(1'-chloroethyl)-5,5-dimethyl-2-(4'-isobutylphenyl)-1,3-dioxane (68.4 g; 0.23 mol), ethylene glycol (92,5 ml), 2,2-dimethyl-1,3-propanediol (36 g) and potassium acetate is heated to 195° C for 2 hours.

10

The mixture is then cooled to 90° C and worked up as described in Example 1a to afford the title product (23.1 g; 0.112 mol). Yield, 48% m.p. 76-77° C.

EXAMPLE 4

15 4'-methoxyphenyl-acetic acid

A mixture of 2-bromoethyl-2-(4'-methoxyphenyl)-4,7-dihydro-1,3-dioxepine (2.99 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and cis-2-butene-1,4-diol (40 ml) is heated to 125° C for 3 hours.

By following the procedure of Example 1a, there is obtained the title product (1.5 g; 0.009 mol). Yield, 90%; m.p. 86-87° C.

20

EXAMPLE 5

2-(5'-bromo-6'-methoxy-2'-naphthyl)-propionic acid

A mixture of 2-(1-bromoethyl)-5,5-dimethyl-2-(5'-bromo-6'-methoxy-2'-naphthyl)-1,3-dioxane (4.58 g; 0.01 mol), potassium acetate (1.2 g; 0.012 mol) and ethylene glycol (40 ml) is heated to 125° C for 24 hours.

25

By following the procedure of Example 1a, there is obtained the title product (2.8 g; 0.009 mol). Yield, 90%; m.p. 161-163° C.

EXAMPLE 6

30 2-(2'-thienyl)-propionic acid

A mixture of 2-(1'-bromoethyl)-5,5-dimethyl-2-(2'-thienyl)-1,3-dioxane (28.7 g; 0.094 mol), ethylene glycol (32.5 g; 0.52 mol), 2,2-dimethyl-1,3-propanediol (13.75 g; 0.13 mol) and potassium acetate (10.25 g; 0.11 mol) is heated to 150° C for 5 hours.

- 5 By following the procedure of Example 1a, there is obtained the title product (11.7 g; 0.075 mol). Yield, 80%; b.p. 98-100° C/0.5 mm Hg.

CLAIMS

1. A process for preparing an alpha-arylalkanoic acid or a salt thereof, which comprises the rearrangement of an alpha-halo-alkyl-arylketal in neutral or weakly alkaline conditions and in the presence of a polar-protic medium and the subsequent hydrolysis of the thus obtained ester in the same reaction medium to afford the corresponding alpha-arylalkanoic acid or a salt thereof.

2. A process according to claim 1, wherein the polar protic medium is consisting of water , an alcohol or a mixture thereof.

3. A process according to claim 2, wherein the alcohol is selected from the group comprising C₁-C₁₂ primary, secondary and tertiary alcohols and C₂-C₁₂ polyhydric alcohols.

4. A process according to claim 1, wherein the neutral or weakly alkaline condition is obtained by adding a buffer, an organic or inorganic weak base or a mixture thereof.

5. A process according to claim 1, wherein the ester is hydrolyzed by means of a base to afford a salt of the corresponding alpha-arylalkanoic acid which, when desired, is treated with an acid to afford the free alpha-arylalkanoic acid.

6. A process according to claim 1, wherein the ester is hydrolyzed by means of an acid to afford the corresponding alpha-arylalkanoic acid.



European Patent
Office

EUROPEAN SEARCH REPORT

0101124

Application number

EP 83 20 1137

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
D, Y	EP-A-0 034 871 (BLASINACHIM) * Claim 1; page 4, lines 6,9; page 5, line 17; pages 22,23, example 2 *	1-3, 5, 6	C 07 C 67/475 C 07 C 69/734 C 07 C 51/00 C 07 C 59/64 C 07 D 333/22 C 07 C 57/30
Y	EP-A-0 035 305 (BLASINACHIM) * Claim 1; page 3, lines 32,35; page 10, example 2 *	1-3, 5, 6	
E	EP-A-0 071 299 (BLASCHIM) * - Claims 1,2,5; pages 4,5, exam- ple 1 *	1-3, 5, 6	
A	GB-A-2 042 543 (MONTEDISON) * Claim 1 *	1	<div>TECHNICAL FIELDS SEARCHED (Int. Cl. 3)</div> C 07 C 59/00 C 07 C 51/00 C 07 C 69/00 C 07 C 67/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 08-11-1983	Examiner KLAG M. J.
<div>CATEGORY OF CITED DOCUMENTS</div> <div> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document </div> <div> T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document </div>			